

REJUVenate®

Ready to Use Wipes

Efficacy Claims & Use Directions



One-Step Disinfectant FOR USE IN SPAS, SALONS & CLINICS

**KEEP OUT OF REACH
OF CHILDREN**

READ S.D.S. BEFORE
USING THE PRODUCT
S.D.S. available online at
www.virox.com

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON CANADA L6H 6R1
1-800-387-7578 | virox.com

Meets OSHA Bloodborne Pathogen Standard for *HBV,*HCV & *HIV-1 | Bactericidal, *Virucidal, Pseudomonacidal, Fungicidal, and Tuberculocidal

For Use In Spas, Salons & Clinics. Just Pull, Wipe & Toss!

This product is a non-woven, disposable wipe pre-saturated with Hydrogen Peroxide disinfectant.

Rejuvenate™ Ready to Use Wipes One-Step Disinfectant is designed for disinfection of hard non-porous surfaces in beauty salons, tanning facilities, spas, nail salons, tattoo parlors, barbershops, and hairstyling establishments.

It is designed for use on hard, non-porous environmental surfaces such as foot baths, foot spas, pedicure foot spa equipment and esthetic tools and equipment.

Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Herpes Simplex Virus, Type 1
Herpes Simplex Virus, Type 2
HIV-1, Strain HTLV-III_B
Human Coronavirus
Influenza A Virus (Strain Hong Kong)
Poliovirus Type 1, Strain Brunhilde
Rhinovirus Type 14
Rotavirus (Strain WA)

Kills Norovirus (Feline Calicivirus as the surrogate) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

BACTERICIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces.

Bactericidal Activity:

Acinetobacter baumannii
Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Shigella dysenteriae
Staphylococcus aureus

Antibiotic-Resistant Bactericidal Activity:

Enterococcus faecalis (Vancomycin Resistant (VRE))
Escherichia coli (Extended Spectrum Beta-Lactamase (ESBL))
Staphylococcus aureus (Community Associated Methicillin Resistant (CA-MRSA)) (NARSA NRS384) (Genotype USA 300)
Staphylococcus aureus (Community Associated Methicillin Resistant (CA-MRSA)) (NARSA NRS123) (Genotype USA 400)
Staphylococcus aureus (Methicillin Resistant (MRSA))

*VIRUCIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces.

Adenovirus type 8
Avian Influenza A virus

TUBERCULOCIDAL: 5 MINUTES

In the presence of 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

†*Mycobacterium bovis* (BCG)

FUNGICIDAL: 10 MINUTES

In the presence of 5% serum load and 10 minute contact time on hard, non-porous environmental surfaces.

Trichophyton interdigitale (Athlete's Foot and Nail Fungus)

NON-FOOD CONTACT SANITIZER: 30 SECONDS

In the presence of 5% serum load and 30 second contact time on hard, non-porous environmental surfaces.

Enterococcus faecalis (VRE)
Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Staphylococcus aureus
Staphylococcus aureus (MRSA)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

For Use as a One-Step Disinfectant:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for Mycobacteria[†]. Use a 10 minute contact time for fungi.
4. Wipe surfaces dry or allow to air dry.

For Use as a Tuberculocide:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 5 minutes. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

To Kill Fungi:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 10 minutes. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.

For use as a Non-Food Contact Sanitizer:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

For use as a disinfectant on hard, non-porous surfaces of foot baths (pedicure foot spa equipment):

1. **Drain:** Drain the water and remove any visible debris.
2. **Clean:** Thoroughly clean all surfaces with soap detergent, then rinse with water and drain.
3. **Apply:** Pull towelettes from canister and wipe all hard non-porous surfaces with the wipe.
4. **To Disinfect:** All surfaces must remain wet for 1 minute. Use a 5 minute contact time for Mycobacteria[†] and a 10 minute contact time for fungi. Use additional towelettes if needed to ensure wet contact time.
5. **Rinse** all surfaces with fresh water.

Note: Change towelettes frequently to avoid re-deposition of soil.

***KILLS HIV AND HCV AND HBV ON pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of Rejuvenate™.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Mycobacteria[†] and 10 minute contact for fungi.

Disposal of Infectious Material: Cleaning materials used that may contain blood and other body fluids must be autoclaved and/or disposed of according to Federal, State, and local regulations for infectious waste disposal.

AHP® and Design are registered trademarks of Diversey Inc.

Si no puede leer en inglés, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto. (If you cannot read English, ask your supervisor to explain the appropriate directions for use before working with this product.)